

Ovarian cryopreservation and transplantation: basic aspects

Kutluk Oktay *, Guvenc G. Karlikaya, Bulent A. Aydin

Department of Obstetrics and Gynecology, New York Methodist Hospital/Cornell University Weill Medical College, 506 6th Street, Brooklyn, NY 11215, USA

Abstract

Animal studies and studies using human ovarian xenografts in immunodeficient mice have shown that ovarian transplantation is feasible. In this manuscript, we present the first and limited experience with ovarian tissue cryopreservation and transplantation in the human. Despite the progress, more basic studies are needed to improve the survival of primordial follicles during and after the cryopreservation process. © 2000 Elsevier Science Ireland Ltd. All rights reserved.

Keywords: Ovary; Transplantation; Cryopreservation; Primordial follicle

1. Introduction

Ovarian tissue banking relies on the principle that the primordial follicles withstand the cryo-toxicity better than the growing follicle (Oktay et al., 1997). A relatively inactive metabolism, lack of zona pellucida, and metaphase spindle grant this privilege to primordial follicle (Oktay et al., 1998a). In addition, smaller cell size allows faster penetration of cryoprotectants in the oocyte. It has been possible to isolate primordial follicles (Oktay et al., 1997) but the only practical way of preserving these follicles at the present is to cryopreserve them in the tissue to perform autotransplantation.

Antral follicles contain oocytes either in prophase-I or metaphase-II stage, and their oocytes are much more liable to cryo-damage (Table 1). In addition to large cell size and zona pellucida, metaphase II oocytes have an additional disadvantage over prophase I oocytes: they have a cell spindle. Studies have shown that even minor variations in ambient temperature can result in damage to the tubulin proteins in the spindle, albeit some of the damage may be reversed by rewarming (Pickering et al., 1990; Gook et al., 1993). The spindle is a dynamic structure composed of microtubules which are continuously being assembled at one end and removed at the other in a treadmilling fashion (George et al., 1996). Although the chromosomes can reassemble

and align along the spindle equator on rewarming the cell, there is a risk of chromosomal loss and aneuploidy occurring at the first maturation division (Oktay and Gosden, 1999). It has also been shown that the cytoskeleton of the antral follicle oocytes is damaged by the cryopreservation process, which could lead to significant changes in the organization and trafficking of molecules and organelles (Vincent and Johnson, 1992). In addition, studies have shown that cryopreservation may result in hardening of zona pellucida, preventing natural fertilization (Oktay and Gosden, 1999).

2. Animal studies

Whole-organ and fetal ovarian tissue cryopreservation and transplantation have been a great success in the rodents (Parrot, 1960; Harp et al., 1994; Cox et al., 1996). However, human ovary is much larger and fibrous, and therefore better animal models were required before the technique could be applied to humans.

Sheep ovary is similar to human ovary, in that; it has a dense fibrous stroma and relatively high primordial follicle density in the ovarian cortex. Gosden and colleagues have performed a two-part study in the sheep. In the first, frozen-banked strips of ovarian cortical pieces were autotransplanted on the infundibulopelvic ligament (Gosden et al., 1994a). Each animal also had a fresh transplant on the opposite site serving as a

* Corresponding author. Fax: +1-718-7803079.

control. A total of 4 months after the transplant, first signs of ovulation were detected. Two pregnancies had occurred, one from a fresh another from a frozen-thawed graft. In the second study, Baird and colleagues have performed autotransplants with frozen tissue in eight sheep and followed them for up to 22 weeks (Baird et al., 1999). All the animals resumed cyclicity and showed hormone production. In that study, baseline levels of FSH were elevated but luteal phase was progesterone measurements were normal. However, serum inhibin-A levels were found to be low in the luteal phase.

3. Studies involving human ovarian tissue

SCID-mice carry a genetic mutation, which results in T- and B-cell immune-deficiency (Bosma et al., 1983). This allows xenografts to revascularize and survive in these animals without being rejected. Roger Gosden has adopted this model for human ovarian xenografting. In his earlier studies, both marmoset and sheep ovarian tissues were transplanted under the kidney capsule, and they grew to antral stages (Gosden et al., 1994b). In the first study using human tissue, Gosden and colleagues cryopreserved ovarian cortical pieces with various cryoprotectants and grafted into SCID mice (Newton et al., 1996). After 18 days, the grafts were removed and primordial follicle counts were obtained. With the exception of glycerol, all cryoprotectants (propanediol, ethylene glycol, DMSO) did well and 44–84% of the follicles survived.

After achieving successful results with short-term xenografting, we performed the two long-term studies

in SCID mice using human tissue (Oktay et al., 1998b; Oktay et al., in press). In the first study, fresh ovarian tissue was used (Oktay et al., 1998b). A 1 mm³ ovarian piece from a 17 year-old patient were grafted under the kidney capsules of hypogonadal SCID mice. During the last 6 weeks of 17-weeks grafting period, one group of animals received FSH stimulation. Estradiol levels were measured at the end of 22 weeks, when animals were sacrificed.

In the FSH treated group, up to 5 mm size antral follicles were found, estradiol levels peaked > 700 pg/ml and the uteri showed clear signs of estrogenization. Because the animals were oophorectomized, the only source of estrogen was human ovarian xenografts.

In the second study, we grafted frozen-thawed human ovarian tissue into SCID mice (Oktay and Karlikaya, 2000). Because these animals were not hypogonadal, no FSH was given. Grafts were recovered 22 weeks later. Histological examination showed follicle growth initiation and an abundant number of healthy primordial follicles remaining. Presumably, because no exogenous FSH was given, further follicle development did not occur. However, this study further strengthened the notion that human primordial follicles survive the triple insult of freeze-thaw-ischemia in long-term transplants.

Finally, to further confirm and quantify the survival of primordial follicles of freezing process, we have performed viability studies on isolated human primordial follicles (Oktay et al., 1997). Ovarian cortical pieces (2 × 2 mm) were cryopreserved with the slow-freeze process (Table 2). After thawing, the tissues were partially digested using collagenase type IA (Sigma, USA) followed by microdissection of primordial follicles.

Table 1
Comparison of available options for banking follicles, oocyte and ovarian tissue

| Cell/tissue type | Size | Advantage | Disadvantage |
|---------------------|-----------------------------|---|---|
| Primordial follicle | 30–50 µm | Least differentiated No ZP ^a No cell spindle | Difficult to isolate Damage to basement membrane Culture not yet possible |
| Preantral follicle | 60–200 µm | Easier to isolate No cell spindle Culture possible | ZP damage More differentiated |
| Prophase I oocyte | 80–100 µm | No spindle Short in vitro maturation | ZP damage Low IVM and fertilization ^b Very few live births |
| Metaphase II oocyte | 80–100 µm | Easy to obtain | ZP damage Cell spindle damage Organelle damage Very few live births |
| Ovarian cortex | 1 × 1 mm to 1 × 3 cm strips | Easy to obtain Preserves stroma Can restore fertility Preventive before cancer treatment | Experimental Risk of cancer cell reseeded |

^a ZP, zona pellucida.

^b IVM, in vitro maturation.

Table 2
Protocol for cryopreserving human ovarian tissue

- (1) Equilibrate 1–3 mm thick slices of ovarian cortex for 30 min on ice in buffered medium containing 1.5 M DMSO (propanediol/ethylene glycole) 10% serum and 0.1 M sucrose
- (2) Load the tissue in cryovials into an automated freezer starting at 0°C and cool at 2°C/min to –7°C
- (3) Soak for 10 min before manual seeding
- (4) Continue to cool at 0.3°C/min to –40°C
- (5) Cool at the faster rate of 10°C/min to –140°C
- (6) Transfer to liquid nitrogen dewar for storage
- (7) Thaw at room temperature for 30 s than at 37°C for 2 min
- (8) Wash tissues stepwise in containing progressively lower concentrations of cryoprotectant media (1.5, 1.0, 0.5, 0 M)

These follicles were then incubated with viability stains. We found that, approximately 70% of follicles survived this process and were viable. Later electron microscopic studies, however, showed that this digestion method might damage the basement membrane of the follicle but the oocyte is rarely affected (Oktay, 1998).

4. Clinical trials of human ovarian tissue cryopreservation and transplantation

Encouraged by these laboratory studies, we have established a human ovarian transplantation project. In the first-phase, only patients aged <35 years were enrolled, and, malignancy cases were excluded.

In the 32 year-old first patient, ovarian tissue was grafted in the forearm into the brachioradialis muscle, akin to the method used for parathyroid gland transplantation. We also grafted tissue in the broad ligament, adjacent to the uterine artery. A total of 4 months after the procedures, ultrasound follow-up indicated that grafts were still intact, and some early antral follicle development was noted in the forearm by high-frequency ultrasound probes. A gradient was detected for estradiol, between the antecubital vein (which the graft drains to) and wrist vein, indicating hormonal production by the graft.

In the second patient, frozen-banked tissue was thawed 8 months after storage, sewn together microscopically to form two large grafts, and transplanted laparoscopically in the ovarian fossa (Oktay and Karlikaya, 2000). Early findings indicated the normalization of serum testosterone levels. Approximately 4 months later, the patient ovulated in response to ovulation induction with human menopausal gonadotropins (Oktay and Karlikaya, 2000).

5. Conclusions

There has been significant progress in ovarian tissue cryopreservation and transplantation in the past 5 years. Now, large-scale human transplantation studies are needed to test the efficacy of this procedure. In addition, research is needed to develop better cryoprotectants and cryopreservation protocols as well as new transplantation techniques to reduce follicle losses that normally occur during the freeze-thaw and grafting process.

References

- Baird, D.T., Webb, R., Campbell, B.K., Harkness, L.M., Gosden, R.G., 1999. Long-term ovarian function in sheep after ovariectomy and transplantation of autografts stored at –196°C. *Endocrinology* 140, 462–471.
- Bosma, G.C., Custer, R.P., Bosma, M.J., 1983. A severe combined deficiency mutation in the mouse. *Nature* 301, 527–530.
- Cox, S.-L., Shaw, J.M., Jenkin, G., 1996. Transplantation of cryopreserved fetal ovarian tissue to adult recipients of mice. *J. Reprod. Fertil.* 107, 315–322.
- George, M.A., Pickering, S.J., Braude, P.R., Johnson, M.H., 1996. The distribution of alpha- and beta-tubulin in fresh and aged human and mouse oocytes exposed to cryoprotectant. *Hum. Reprod.* 2, 445–456.
- Gook, D.A., Osborn, S.M., Johnston, W.I.H., 1993. Cryopreservation of mouse and human oocytes using 1,2-propanediol and the configuration of the meiotic spindle. *Hum. Reprod.* 8, 1101–1109.
- Gosden, R.G., Baird, D.T., Wade, J.C., Webb, R., 1994a. Restoration of fertility to oophorectomized sheep by ovarian autografts stored at –196°C. *Hum. Reprod.* 9, 597–603.
- Gosden, R.G., Boulton, M.I., Grant, K., et al., 1994b. Follicular development of ovarian xenografts in SCID mice. *J. Reprod. Fertil.* 101, 619–623.
- Harp, R., Leibach, J., Black, J., Keldahl, C., Karow, A., 1994. Cryopreservation of murine ovarian tissue. *Cryobiology* 31, 336–343.
- Newton, H., Aubard, Y., Rutherford, A., Sharma, V., Gosden, R., 1996. Low temperature storage and grafting of human ovarian tissue. *Hum. Reprod.* 11, 1487–1491.
- Oktay, K., 1998. New horizons in assisted reproduction. *Assist. Reprod. Rev.* 8, 51–54.
- Oktay, K., Gosden, R.G., 1999. Cryopreservation of human oocytes and ovarian tissue. In: Brinsden, P.R. (Ed.), *A Textbook of in vitro Fertilization and Assisted Reproduction*. Parthenon Publishing, London, UK.
- Oktay, K., Newton, H., Gosden, R.G. Transplantation of banked human ovarian tissue results in follicle growth initiation in SCID mice. *Fertil. Steril.* in press.
- Oktay, K., Nugent, D., Newton, H., Salha, O., Gosden, R.G., 1997. Isolation and characterization of primordial follicles from fresh and cryopreserved human ovarian tissue. *Fertil. Steril.* 67, 481–486.
- Oktay, K., Newton, H., Aubard, Y., Gosden, R.G., 1998a. Cryopreservation of human oocytes and ovarian tissue: an emerging technology? *Fertil. Steril.* 69, 1–7.
- Oktay, K., Newton, H., Mullan, J., Gosden, R.G., 1998b. Development of human primordial follicles to antral stages in SCID/hpg mice stimulated with follicle stimulating hormone. *Hum. Reprod.* 13, 1133–1138.

- Oktay, K., Karlikaya, G., 2000. Ovarian function after transplantation of frozen, banked autologous ovarian tissue. *N. Engl. J. Med.* 342, 1919.
- Parrot, D.M., 1960. The fertility of mice with orthotopic grafts derived from frozen tissue. *J. Reprod. Fertil.* 1, 230–241.
- Pickering, S.J., Braude, P.R., Johnson, M.H., Cant, R.M., Currie, R.M., 1990. Transient cooling to room temperature can cause irreversible disruption of the meiotic spindle in the human oocyte. *Fertil. Steril.* 54, 102–108.
- Vincent, C., Johnson, M.H., 1992. Cooling, cryoprotectants, and the cytoskeleton of the mammalian oocyte. *Oxf. Rev. Reprod. Biol.* 14, 73–100.

AUTOMATED REQUEST TO THE BRITISH LIBRARY DOCUMENT SUPPLY CENTRE
AS-2000/229 PHOTO

B COPY

EMAIL REQUEST

R.D.
05-Mar-02180

P.D.
05-Mar-02193

Customer Code No.
50-0039

Shelfmark
5900.760000

MOL CELL ENDOCRINOL
OVARIAN CRYOPRESERVATION AND
TRANSPLANTATION...

2000 27 169 1-2 105-108

OKTAY K, AYDIN BA

SEE ESTAR

OS FLOOR 4-3
MOLECULAR AND CELLULAR ENDOCRINO..
SEE ALSO SERIAL RECORDS KCS/SE
D1 S1 FA89-E (31) 2000

COPY



27771821

598



27771821

for RMS
use only

ADILNA SANOVEL ILAC SANAYI TIC Aa
ATTN: DR TIJEN ACARKAN
BUYUKDERE CADDESI, DEREBOYU SOKAK
ZAGRA IO MERKEZI C BLOK
80670 MEDIKAL PAZARLAMA, ISTANBUL
TURKEY

International Loan, Return Airmail within 4
weeks of date of receipt unless recalled earlier.

Request Ref. No.
AS-2000/229 PHOTO

If no other library indicated please return loan to:-
The British Library Document Supply Centre, Boston Spa,
Wetherby, West Yorkshire, United Kingdom, LS23 7BQ

THE BRITISH LIBRARY



This document
has been supplied
by, or on behalf of,
The British Library
Document Supply
Centre,
Boston Spa,
Wetherby,
West Yorkshire
LS23 7BQ
United Kingdom

WARNING:
Further copying of
this document
(including storage in
any medium by
electronic means),
other than that
allowed under the
copyright law, is not
permitted without the
permission of the
copyright owner or
an authorised
licensing body.